






# - Radon causes lung cancer even BELOW EPA's radon action level of 4 pCi/L (150 Bq/m³) - North American, European, and Chinese studies report 8 to 16% increase in lung cancer 2.7 pCi/L (100 Bq/m³) - MOST radon-induced lung cancer occurs below EPA's action level - Protracted radon exposure increases the risk of all types of lung cancer - Prevention and mitigation methods reduce the risk - Susceptibility to radon-induced lung cancer varies (e.g., smoking, genetics) - Protracted radon exposure appears to have adverse health effects beyond lung cancer (e.g., leukemia, stomach and liver cancer) (Field, RW 2008 "Testimony Before the President's Cancer Panel," Charleston, SC [December 4]) Radon Programs KANSAS STATE UNIVERSITY




























# **EPA Has Defined Radon Zones**


Each of 3100 counties in the U.S. classified as:


**Zone 1** - expect 4.0 pCi/L or greater


Zone 2 - expect 2.0 to 4.0 pCi/L


Zone 3 - expect 2.0 pCi/L or less


Zone designations based on five factors: indoor radon measurements, geology, aerial radioactivity, soil parameters, and foundation type.






















# **Common Testing Durations**

### How long does radon testing take?

- · Long Term greater than 90 day exposure
  - No behavioral restrictions on building operation
  - Reflects building annualized radon concentrations
  - Short Term typically 2 7 day exposure
    - Short Term cannot be less than 2 days
    - · Closed building conditions required
  - Reflects building winter radon concentrations 94% of the time Short Term tests provide the same mitigation decision as a Long Term test











### Radon Mitigation


- Active Soil Depressurization (ASD) is the most common approach
  - Employs a method for creating a vacuum beneath the foundation greater in strength than the vacuum applied to the soil by the building
  - Caulking and sealing has not been shown to be an effective stand-alone radon reduction process
    - However extensive sealing can radically increase the efficiency of ASD systems
- Ventilation approaches have proven more costly and less effective





























We're From the Government, We're Here to Help

National Radon Program Services

www.sosradon.org

Kansas Radon Program

www.kansasradonprogram.org

K-State Radon Training

https://radoncourses.com/

K-State Radon Chamber

www.ksuradonchamber.org

Radon Programs

KANSAS STATE

UNIVERSITY



# KANSAS RADON PROGRAM

Kansas Radon Program (KRP)


- https://kansasradonprogram.org/home
- •800.693.5343
- State-wide Radon Activities
  - Public education and technical assistance for Kansas residents
  - Professional education for radon stakeholders
  - Outreach activities to radon stakeholders


Radon Programs KANSAS STATE



K-State Radon Training Options

- https://radoncourses.com/
- In-person, Webinar & Online Self-Paced
  - Full measurement/mitigation courses
  - Professional Continuing Education
  - Contract training for organizations
  - EPA Regional Stakeholders meetings





# Contact Info

### Brian Hanson

- 785.532.6026 (general office)
- 785.532.4996 (direct)
- bhanson@ksu.edu

### NRPS & KRP Contact

• radon@ksu.edu

KSU Radon Training

• <u>radoncourse@ksu.edu</u>

KSU Radon Chamber

• radonchamber@ksu.edu