

Habitat Model for Species: **Yellow-throated Vireo**

Alpha Code: YTVI **Status: Breeding**

[Distribution Map](#)

[Habitat Map](#)

Landcover Category

0 - Comments

[KS GAP]
James, 1971

Habitat Restrictions

Map 30 m edge of listed habitats.

James, 1976

James, 1978
James, 1979

Shackford, 1992

Anderson and Robbins, 1981
Whitcomb et al., 1981 territories found only in forest islands >70 ha.
Bent, 1950

01 - Maple-Basswood Forest

[KS GAP] 30 m edge.
Ambuel and Temple, 1983
James, 1979
Bent, 1950
Rodewald and James, 1996 edge

02 - Oak-Hickory Forest

[KS GAP] 30 m edge.
Kahl et al., 1985
Ambuel and Temple, 1983
James, 1979
Bent, 1950
Rodewald and James, 1996 edge

05 - Ash-Elm-Hackberry Floodplain Forest

[KS GAP] 30 m edge.
Zimmerman and Tatschl, 1975
Rodewald and James, 1996 edge

08 - Bur Oak Floodplain Woodland

[KS GAP] 30 m edge.
Kahl et al., 1985
Rodewald and James, 1996 edge

50 - Deciduous Forest-Mined Land

[KS GAP] 30 m edge.
Rodewald and James, 1996 edge

51 - Maple Floodplain Forest

[KS GAP] 30 m edge.
Bent, 1950
Rodewald and James, 1996 edge

Comments

Order of vireos listed from open-country to shaded forest: Bell's, Warbling, White-eyed, Yellow-throated, and Red-eyed. Listed from decreasing shrub density: White-eyed, Red-eyed, Bell's, Warbling, and Yellow-throated. Require taller trees than others.

Over half of the territories has oak present. Preference for completely dead trees or trees such as oaks and aspens which often have large numbers of dead lower limbs.

"open deciduous forest edge"
Tall mature deciduous forest with widely spaced trees, but often with little or no understory. Nested in thicket of close-set hackberry, soapberry and cottonwood trees near a small stream in western Oklahoma (outside normal range).
found only in 4,000+ ha woodlands


may be found on the edges of woodlands, in groves and in open stands of oaks, maples, and other hardwood trees, but seldom in the dense forests.

Reference List


1. Ambuel, B. and S. A. Temple. 1983. Area-dependent changes in the bird communities and vegetation of southern Wisconsin forests. *Ecology* 64(5):1057-1068.
2. Anderson, S. H. and C. S. Robbins. 1981. Habitat size and bird community management. *Transactions of the North American Wildlife and Natural Resources Conference* 46:511-520.

3. Bent, A. C. 1950. Life Histories of North American Wagtails, Shrikes, Vireos, and their allies. United States National Museum Bulletin. No. 197.
4. James, F. C. 1971. Ordination of habitat relationships among breeding birds. *Wilson Bulletin* 83(3):215-236.
5. James, R. D. 1976. Foraging behavior and habitat selection of three species of vireos in southern Ontario. *Wilson Bulletin* 88(1):62-75.
6. _____. 1978. Pairing and nest site selection in Solitary and Yellow-throated Vireos with a description of a ritualized nest building display. *Canadian Journal of Zoology* 56:1163-1169.
7. _____. 1979. The comparative foraging behaviour of Yellow-throated Vireos and Solitary Vireos: the effect of habitat and sympatry. Pages 137-163 *in* J. G. Dickson, R. N. Conner, R. R. Fleet, J. C. Kroll, and J. A. Jackson, editors. *The role of insectivorous birds in forest ecosystems*. Academic Press, New York, New York, USA.
8. Kahl, R. B., T. S. Baskett, J. A. Ellis, and J. N. Burroughs. 1985. Characteristics of summer habitats of selected nongame birds in Missouri. University of Missouri Agriculture Experiment Station Research Bulletin 1056. University of Missouri, Columbia, Missouri, USA.
9. Rodewald, P. G. and R. D. James. 1996. Yellow-throated Vireo. A. Poole and F. Gill, editors. *The Birds of North America*. No. 247. The American Ornithologists' Union and The Academy of Natural Sciences, Philadelphia, Pennsylvania, USA.
10. Shackford, J. S. 1992. Yellow-throated Vireo nest in Cimarron County, Oklahoma. *Bulletin of the Oklahoma Ornithological Society* 25(2):15-16.
11. Whitcomb, R. F., C. S. Robbins, J. F. Lynch, B. L. Whitcomb, M. K. Klimkiewicz, and D. Bystrak. 1981. Effects of forest fragmentation on avifauna of the Eastern Deciduous Forest. Pages 126-205 *in* R. L. Burgess and D. M. Sharpe, editors. *Forest island dynamics in man-dominated landscapes*. Springer-Verlag, New York, New York, USA.
12. Zimmerman, J. L. and J. L. Tatschl. 1975. Floodplain birds of Weston Bend, Missouri River. *Wilson Bulletin* 87(2):196-208.


Yellow-throated Vireo





Kansas Breeding Bird Atlas


 Confirmed Breeding

 Probable Breeding

 Possible Breeding

 Breeding Bird Survey



 *Birds of Kansas,*
Horned Lark

 Range from Johnsgard, 1998
The Birds of Nebraska

Habitat

All shaded counties will be included in the possible distribution of the species.

Yellow-throated Vireo



* 30 m edge