

Habitat Model for Species: Five-lined Skink

Eumeces fasciatus

[Distribution Map](#)
[Habitat Map](#)

Landcover Category

0 - Comments

Fitch and von Achen, 1977
Webb, 1970
Clarke, 1965

Fitch, 1954
Burt and Hoyle, 1935
Clarke, 1958

Hall and Smith, 1947
Trowbridge, 1937
Fitch, 1958
Hecnar, 1994

Seburn, 1993

[#Reviewer]

Habitat Restrictions

Comments

No useful habitat information
Usually found in woodlands or shaded areas, usually on the ground. May be found in grassy fields where rocks or other materials form protective cover.

No useful habitat information

No useful habitat information

Also found occasionally in buckbrush-sumac community, an intermediate ecological area between the hillside forest and the prairies.

Distribution only

No useful habitat information

No useful habitat information

Study conducted in Canada. Habitat information not used in model.

Study conducted in Canada. Habitat information not used in model.

Fitch Reservation: watched it shrink from abundance to scarcity. I would define its habitat as open deciduous woodland with loose rocks. Trees species include chestnut oak, American elm, hackberry, and walnut.

Busby: In my experience, found in almost any upland forest and woodland with rocky substrate or other type of structure on ground.

Busby: In my experience, found in almost any upland forest and woodland with rocky substrate or other type of structure on ground.

Busby: In my experience, found in almost any upland forest and woodland with rocky substrate or other type of structure on ground.

01 - Maple-Basswood Forest

[#Reviewer]

02 - Oak-Hickory Forest

Clarke, 1958
Gloyd, 1928
Fitch, 1963
Brumwell, 1951
[#Reviewer]

03 - Post Oak-Blackjack Oak Forest

Brumwell, 1951
[#Reviewer]

04 - Pecan Floodplain Forest

Fitch, 1963

05 - Ash-Elm-Hackberry Floodplain Forest

Gloyd, 1928
Fitch, 1963

07 - Mixed Oak Floodplain Forest

Fitch, 1963

08 - Bur Oak Floodplain Woodland

Gloyd, 1928
Fitch, 1963

09 - Mixed Oak Ravine Woodland

Clarke, 1958
Fitch, 1963

10 - Post Oak-Blackjack Oak Woodland

[#Reviewer]

Busby: In my experience, found in almost any upland forest and woodland with rocky substrate or other type of structure on ground.

50 - Deciduous Forest-Mined Land

Clarke, 1958

[#Reviewer]

Busby: In my experience, found in almost any upland forest and woodland with rocky substrate or other type of structure on ground.

55 - Deciduous Woodland

[#Reviewer]

Busby: In my experience, found in almost any upland forest and woodland with rocky substrate or other type of structure on ground.

91 - Ozark Sandstone Glade / Prairie - not mapped

[#Reviewer]

Rundquist: add

92 - Ozark Limestone Glade - not mapped


[#Reviewer]


Rundquist: add

Reference List

1. Brumwell, M. J. 1951. An ecological survey of Fort Leavenworth Military Reservation. *American Midland Naturalist* 45(1):187-231.
2. Burt, C. E. and W. L. Hoyle. 1935. Additional records of the reptiles of the central prairie region of the United States. *Transactions of the Kansas Academy of Science* 37:193-216.
3. Clarke, R. F. 1958. An ecological study of the reptiles and amphibians in Osage County, Kansas. *Emporia State Research Studies, Kansas State Teachers College* 7(1):1-52.
4. _____. 1965. Lizards in Kansas. *Kansas School Naturalist* 11(4):1-16.
5. Fitch, H. S. 1954. Life history and ecology of the five-lined skink, *Eumeces fasciatus*. University of Kansas Publications, Museum of Natural History 8(1):1-156.
6. _____. 1958. Home ranges, territories, and seasonal movements of vertebrates of the Natural History Museum. University of Kansas Publications, Museum of Natural History 11(3):63-326.
7. _____. 1963. Natural history of the black rat snake (*Elaphe o. obsoleta*) in Kansas. *Copeia* 4:649-658.
8. Fitch, H. S. and P. L. von Achen. 1977. Spatial relationships and seasonality in the skinks *Eumeces fasciatus* and *Scincella laterale* in northeastern Kansas. *Herpetologica* 33(3):303-313.
9. Gloyd, H. K. 1928. The amphibians and reptiles of Franklin County, Kansas. *Transactions of the Kansas Academy of Science* 31:115-141.
10. Hall, H. H. and H. M. Smith. 1947. Selected records of reptiles and amphibians from southeast Kansas. *Transactions of the Kansas Academy of Science* 49(4):447-454.
11. Hecnar, S. J. 1994. Nest distribution, site selection, and brooding in the five-lined skink (*Eumeces fasciatus*). *Canadian Journal of Zoology* 72:1510-1516.
12. Seburn, C. N. L. 1993. Spatial distribution and microhabitat use in the five-lined skink (*Eumeces fasciatus*). *Canadian Journal of Zoology* 71:445-450.
13. Trowbridge, A. H. 1937. Ecological observations on amphibians and reptiles collected in southeastern Oklahoma during the summer of 1934. *American Midland Naturalist* 18(2):285-303.
14. Webb, R. G. 1970. *Reptiles of Oklahoma*. University of Oklahoma Press, Norman, Oklahoma, USA.

Five-lined Skink



 Range from Collins, 1993

Five-lined Skink

