

Habitat Model for Species: Northern Prairie Skink

Eumeces septentrionalis septentrionalis

[Distribution Map](#)
[Habitat Map](#)

Landcover Category

0 - Comments

Fleharty and Ittner, 1967
Webb, 1970
Clarke, 1965

Stains and Ozment, 1962
Rundquist et al., 1978
Burt and Hoyle, 1935
Clarke, 1956
Clarke, 1955
Gehlbach and Collette, 1959
Breckenridge, 1943
Fitch, 1958
[#KS GAP]

[#Reviewer]

Habitat Restrictions

Comments

Distribution only
No useful habitat information.
Most frequently are found in open areas, particularly in rocky places on grassy prairie hills.

Distribution only
Distribution only
Distribution only
Distribution only
No useful habitat information
Distribution in Nebraska only.
No useful habitat information
No useful habitat information
The draft model for Northern and Southern Prairie Skinks were done together as one model.
Rundquist: This species is always associated with rock outcrops.
Fitch Reservation: Formerly occurred here but was very scarce and localized. I have seen only a few in mowed grass to create a lawn. Absent from surrounding mixed

05 - Ash-Elm-Hackberry Floodplain Forest

[#Reviewer]

Busby: I have found this species several times in Ash-Elm-Hackberry Floodplain Forest in Saline County.

17 - Tallgrass Prairie

Clarke, 1955

55 - Deciduous Woodland


[#Reviewer]


Rundquist: add

Reference List

1. Breckenridge, W. J. 1943. The life history of the black-banded skink *Eumeces septentrionalis* (Baird). American Midland Naturalist 29:591-601.
2. Burt, C. E. and W. L. Hoyle. 1935. Additional records of the reptiles of the central prairie region of the United States. Transactions of the Kansas Academy of Science 37:193-216.
3. Clarke, R. 1955. Observations on *Eumeces s septentrionalis* in Kansas. Herpetologica 11(3):161-164.
4. Clarke, R. F. 1956. Distributional notes on some amphibians and reptiles of Kansas. Transactions of the Kansas Academy of Science 59(2):213-219.
5. _____. 1965. Lizards in Kansas. Kansas School Naturalist 11(4):1-16.
6. Fitch, H. S. 1958. Home ranges, territories, and seasonal movements of vertebrates of the Natural History Museum. University of Kansas Publications, Museum of Natural History 11(3):63-326.
7. Fleharty, E. D. and D. R. Ittner. 1967. Additional locality records for some Kansas Herptiles. Southwestern Naturalist 12(2):199-200.
8. Gehlbach, F. R. and B. B. Collette. 1959. Distributional and biological notes on the Nebraska herpetofauna. Herpetologica 15:141-143.
9. Rundquist, E. M., E. Stegall, D. Grow, and P. Gray. 1978. New herpetological records from Kansas. Transactions of the Kansas Academy of Science 81(1):73-77.
10. Stains, H. J. and J. Ozment. 1962. A record of the brown skink (*Scincella laterale*) and prairie skink (*Eumeces septentrionalis*) from Barber County, Kansas. Transactions of the Kansas Academy of Science 65(2):143.
11. Webb, R. G. 1970. Reptiles of Oklahoma. University of Oklahoma Press, Norman, Oklahoma, USA.

Northern Prairie Skink



 Range from Collins, 1993

Northern Prairie Skink

