Getting Unstuck

Student Workbook



KSU Counseling and Psychological Services

785-532-6927

www.k-state.edu/counseling/

Adapted from Cal Poly Getting Unstuck Program

Table of Contents

In Session Worksheets	Page 3
Session 1: Common Depressive Symptoms	Pages 4
Common Schemas	Pages 5-6
Common Depressive Symptoms	Page 7
Cross-Sectional Formulation (example)	Page 8
Cross-Sectional Formulation (blank)	Page 9
Feelings Wheel	Page 10
Sleep Hygiene and Exercise Tip	Page 11
Session 1 Homework Assignment	<u>Page 12</u>
Cross-Sectional Formulation (blank)	Pages 13 -16
Session 2: Unhelpful Thoughts and Behaviors	Page 17
How To Do a "Good" Depression	Page 18
Looking at Thoughts	Page 19
Unhelpful Thinking Styles	Page 20
Unhelpful Behaviors	Page 21
<u>Session 2 Homework Assignments</u>	<u>Page 22</u>
Homework: Maintaining Depressed Mood	Page 23
Cross-Sectional Formulation and Looking at Thoughts (blank)	Page 24 - 29
Session 3: Modifying Thoughts and Behaviors	Page 30
Self-Affirmations	Page 31
Alternative Response Worksheet (example)	Page 32
Alternative Response Worksheet (blank)	Page 33
Challenging the Unhelpful Thinking Styles	Page 34
If you're having trouble, ask yourself these questions	Page 35
Feelings Wheel	Page 36
Other Helpful Thought Tips	Page 37
Helpful Behavioral Tips	Page 38
Other Helpful Behavioral Tips for Managing Depression	Page 39
Putting it All Together: My Personalized Plan	Page 40
Appendix	Page 42 - 43

IN SESSION WORKSHEETS

SESSION 1: DEPRESSION 101

Common Depressive Symptoms

Self-criticism	Reduced focus/Concentration	Low mood or persistent sadness	Fatigue or low energy		
Withdrawing from friends, family, pets	Suicidal thoughts or preoccupation with death	Crying Spells	Suicide attempts		
Changes to sex drive	Poor hygiene	Anger or irritability	"Self-medicating" with alcohol or other drugs		
Changes to eating or weight	Worry about something awful happening	Guilt	Appetite disruption		
Unexplained aches and pains	Numbness or feelings of emptiness	Increased or reduced sleep	Gastrointestinal problems		
Physical restlessness or slowing	"Heaviness" in the body	Hopelessness	Fear of rejection		
	Loss of Pleasure for things you used to enjoy	Inability to make decisions			

Common Schemas

- **Abandonment**: The belief and expectation that others are unreliable and will eventually leave. The belief that relationships are fragile, loss is inevitable, and you will ultimately wind up alone.
- **Approval-Seeking/Recognition-Seeking:** The sense that approval, attention, and recognition from others are far more important than genuine self-expression or being true to yourself.
- **Defectiveness/Shame:** The belief that you are flawed, unlovable, or, damaged and would be rejected by significant others if exposed. May involve hypersensitivity to criticism, insecurity around others, or a sense of shame regarding perceived flaws.
- **Dependence/Incompetence:** The belief that your judgement is questionable and you are incompetent. This leads to feeling like you are unable to handle day-to-day responsibilities (i.e., taking care of yourself or making good choices) without considerable help from others.
- **Emotional Deprivation:** Expectation that your primary emotional needs, including nurturing, protection, and empathy, will never be met.
- **Emotional Inhibition:** The belief that you must control your self-expression or others will reject or criticize you.
- Enmeshment/Undeveloped Self: The belief that you do not have an individual identity separate from one or more significant others, resulting in unhelpful levels of emotional involvement and closeness. May also include feelings of smothering.
- Entitlement/Grandiosity: The belief that you are special or more important than other people and entitled to special rights and privileges, even though it may have a negative effect on others. May include an exaggerated focus on superiority in order to achieve power or control.


- **Failure:** The belief that you have failed, will inevitably fail, or are fundamentally inadequate in areas of achievement (i.e., school, career, sports, etc.).
- **Insufficient Self Control/Self-Discipline:** Difficulty or refusal to exercise self-control and frustration tolerance in order to achieve personal goals, or to restrain the excessive expression of emotions and impulses.
- **Mistrust/Abuse:** The belief that others will hurt, abuse, humiliate, cheat, or manipulate you. Usually involves the belief that the harm is intentional or due to extreme negligence.
- **Negativity/Pessimism:** A pervasive belief that the negative aspects of life outweigh positive or optimistic aspects. The expectation that things will eventually go seriously wrong or that things that seem to be going well will ultimately fall apart.
- **Punitiveness:** The belief that people should be harshly punished for their mistakes or shortcomings.
- **Self-Sacrifice:** The belief that **you should** focus on meeting the needs of others at the expense of your own needs, to the point that it is excessive and harmful.
- **Social Isolation:** The belief that you are isolated from the rest of the world, different from other people, and/or alienated from a community.
- **Subjugation:** Excessive surrendering of control to others because you feel coerced, usually to avoid anger, retaliation, or abandonment.
- **Unrelenting Standards:** The belief that you need to be the best, constantly striving for perfection or trying to avoid mistakes.
- **Vulnerability:** Belief that the world is a dangerous place and that imminent catastrophe will strike at any time and that you will be unable to prevent it.


Common Depressive Symptoms

Behaviors	Thoughts	Emotions	Physical Sensations		
Increased or reduced sleep	Reduced focus/concentration	Low Mood or persistent sadness	Fatigue or low energy		
Withdrawing from friends, family, pets	Suicidal thoughts or preoccupation with death	Loss of Pleasure for things you used to enjoy	Unexplained aches and pains		
"self- medicating" with alcohol or other drugs	Self-criticism	Anger or irritability	"heaviness" in the body		
Changes to eating or weight	Worry about something awful happening	Guilt	Appetite disruption		
Suicide attempts	Fear of rejection	Numbness or feelings of emptiness	Gastrointestinal problems		
Physical restlessness or slowing	Inability to make decisions	Hopelessness	Changes to sex drive		
Poor hygiene					
Crying Spells					


Situation (When? Where? What? With whom?) I have a group assignment due for my class. I asked a fellow classmate to be my partner, but she declined.

Schemas: failure, abandonment, defectiveness, social isolation





Feelings Wheel



Sleep Hygiene



- 1. **Get regular.** Go to bed and get up at more or less the same time every day, even on weekends and days off!
- 2. **Get up and try again.** Try to go to sleep only when tired. If you haven't been able to get to sleep after about 30 minutes, get up and do something calming (not stimulating) until you feel sleepy, then return to bed and try again.
- 3. Avoid caffeine and nicotine. Avoid consuming any caffeine (coffee, tea, soda, chocolate) or nicotine (cigarettes) for at least 4-6 hours before going to bed.



These act as stimulants and interfere with falling asleep.

- 4. **Avoid alcohol**. Avoid alcohol for at least 4-6 hours before bed because it interrupts the quality of sleep.
- 5. **Bed is for sleeping.** Try not to use your bed for anything other than sleeping and sex, so that your body comes to associate bed with sleep.
- 6. **Electronics curfew.** Don't use back-lit electronics 60 minutes prior to bed, as the artificial light inhibits hormones and neurons that promote sleep.

- 7. **No naps.** Avoid taking naps during the day. If you can't make it through the day without a nap, make sure it is for less than an hour and before 3pm.
- 8. **Sleep rituals**. Develop rituals to remind your body that it is time to sleep, like relaxing stretches or breathing exercises for 15 minutes before bed.



- 9. No clock-watching. Checking the clock during the night can wake you up and reinforces negative thoughts such as, "Oh no, look how late it is, I'll never get to sleep."
- 10. **The right space.** Make your bed and bedroom quiet and comfortable for sleeping. An eye mask and earplugs may help block out light and noise.
- 11. **Keep daytime routine the same.** Even if you have a bad night's sleep, it is important that you try to keep your daytime activities the same as you had planned. That is, don't avoid activities because you feel tired. This can reinforce the insomnia.

Exercise Tips

- 1. Find an enjoyable activity. Exercise doesn't have to be boring. Choose a pleasurable activity, like playing badminton or doing yoga. Aim for fun, not more work.
- 2. Start small. Commit to 10 minutes of exercise a day or add exercise to your daily routine (like walking to school or doing push-ups in your room).
- **3. Get outside.** The sun provides a mood "pick me up" of its own, producing serotonin in the brain. Take a walk outside or go swimming.



- **4.** Schedule it in. It's easy to skip exercise when we don't plan. Put it in your phone as part of your daily to-do's and celebrate when you check it off.
- **5. Mix it up.** To avoid feeling bored with exercise, try a number of different activities.

- 6. **Team up.** Depression can be isolating. Ask others to do team activities or find an exercise buddy for accountability and to increase social interaction.
- 7. Minimize equipment. Equipment can be expensive. Identify activities that don't require you to have equipment or facilities, like walking, running, or dancing.





- 8. Follow your energy. If your energy fluctuates throughout the day, try to plan to exercise when your energy is at its peak. Alternatively, exercise when feeling sluggish for an energy boost.
- **9. Set goals.** Achieving goals improves mood and selfesteem. Set specific achievable exercise goals and reward yourself when you accomplish them.


Session 1 Homework Assignments


Homework 1: Complete at least one cross sectional formulation for review next week (multiple blank copies provided starting on page 13)

Homework 2: Implement one sleep hygiene and one exercise tip over the next week.









SESSION 2: UNHELPFUL THOUGHTS AND BEHAVIORS

How To Do a "Good" Depression

- Stay still, don't do anything.
- Stay in bed if you can; if not, sit in the same chair or lay on the couch.



- Watch excessive amounts of TV or go online for hours, particularly viewing social media.
- Sleep during the day and don't sleep at night.
- Focus on the past, your fears, faults and resentments.



- Imagine the future will be the same or worse than the past or present.
- Isolate; avoid other people, including friends and family.

- If you can't avoid other people, try to talk to the same person or few people.
- If you do talk to people, talk about the same topic, usually how depressed or unhappy you are.
- Eat poorly; overeat or stop eating. Eat predominantly junk food, sugar, and carbs.



- Don't pursue hobbies, passions, or interests.
- Drink alcohol, smoke cigarettes, and/or use other drugs.
- Don't ask for help.

Adapted from Bill O'Hanlon's "Do One Thing Different"


Looking at Thoughts

Cognitive (What went through your mind when you felt depressed? What did that say or mean about you or the situation?)

Schemas:

- Abandonment
- Approval-Seeking/ Recognition-Seeking
- Defectiveness/Shame
- Dependence/Incompetence
- Emotional Deprivation
- Emotional Inhibition
- Enmeshment/Undeveloped Self
- Entitlement/Grandiosity
- Failure

- Insufficient Self Control/Self-Discipline
- Mistrust/Abuse
- Negativity/Pessimism
- Punitiveness
- Self-Sacrifice
- Social Isolation
- Subjugation
- Unrelenting Standards
- Vulnerability



Unhelpful Thinking Styles

Unhelpful Behaviors

Get up at different times every morning	Go to bed at different times every night	Miss class	Don't do a homework assignment	Wait until the last minute to get something done	Stop washing your hair Over eat
Stop brushing your teeth	Skip a test	Stop going to team or club meeting	Ignore or decline invites to hang out	decline invites to	
Eat lots of junk food	Binge drink	Smoke cigarettes	Stop talking to/activelyStop talking to/activelyavoid your friendsavoid your family		Don't seek help when you are struggling
Stay inside all the time	Play video games for hours	Constantly monitor social media	Stay in your Stay in your bed room; don leave		Stop talking to your roommate
Stop completing tasks	Stop going to work	Stop exercising	Watch a lot of TV	Only eat one meal a day	Argue with other people
Stop doing your hobbies	Only focus on the negative	Complain a lot	Use recreational drugs	Sleep all day	Don't move
Only listen to music that makes you sad, angry or upset	Stop smiling and laughing	Stop going to church, meditating or praying	Let your room get really disorganized	Self-harm	Smoke marijuana

Session 2 Homework Assignments

Homework 1: Complete "Maintaining Depressed Mood" worksheet (page 23)

Homework 2: Complete at least one Simple Cross Sectional Formulation worksheet for review next week (multiple blank copies provided starting on page 24)

Maintaining Depressed Mood

Your task is to be a curious investigator or scientist in the study of you for at least two days this week. What are your thought and behavior patterns that perpetuate feeling depressed? What perpetuates feeling okay or good? What is not working and what is working?

Things I think and do that maintain my depressed mood:

Examples of thoughts: Magnification, disqualifying the positive, emotional reasoning

<u>Examples of behaviors:</u> Skip meals, stop exercising, stay up late, stop showering, miss class, avoid friends


Day	Thoughts	Behaviors
1		
2		
(Sat or Sun)		

Things I think and do that help me feel okay, good, and/or accomplished:

<u>Examples of thoughts:</u> Compliment yourself, give yourself credit for doing something, talk back to your negative thoughts, remind yourself to be gentle to yourself, think about a good memory

<u>Examples of behaviors</u>: Read a book for fun, go to a club meeting, brush my teeth at night, go to the library to study, clean/organize my room, eat breakfast, take vitamins, call a friend, go to a coffee shop, do a chore

Day	Thoughts	Behaviors
1		
2		
(Sat or Sun)		
()		




Looking at Thoughts

Cognitive (What went through your mind when you felt depressed? What did that say or mean about you or the situation?)

Schemas:

- Abandonment
- Approval-Seeking/ Recognition-Seeking
- Defectiveness/Shame
- Dependence/Incompetence
- Emotional Deprivation
- Emotional Inhibition
- Enmeshment/Undeveloped Self
- Entitlement/Grandiosity
- Failure

- Insufficient Self Control/Self-Discipline
- Mistrust/Abuse
- Negativity/Pessimism
- Punitiveness
- Self-Sacrifice
- Social Isolation
- Subjugation
- Unrelenting Standards
- Vulnerability




Looking at Thoughts

Cognitive (What went through your mind when you felt depressed? What did that say or mean about you or the situation?)

Schemas:

- Abandonment
- Approval-Seeking/ Recognition-Seeking
- Defectiveness/Shame
- Dependence/Incompetence
- Emotional Deprivation
- Emotional Inhibition
- Enmeshment/Undeveloped Self
- Entitlement/Grandiosity
- Failure

- Insufficient Self Control/Self-Discipline
- Mistrust/Abuse
- Negativity/Pessimism
- Punitiveness
- Self-Sacrifice
- Social Isolation
- Subjugation
- Unrelenting Standards
- Vulnerability



Looking at Thoughts

Cognitive (What went through your mind when you felt depressed? What did that say or mean about you or the situation?)

Schemas:

- Abandonment
- Approval-Seeking/ Recognition-Seeking
- Defectiveness/Shame
- Dependence/Incompetence
- Emotional Deprivation
- Emotional Inhibition
- Enmeshment/Undeveloped Self
- Entitlement/Grandiosity
- Failure

- Insufficient Self Control/Self-Discipline
- Mistrust/Abuse
- Negativity/Pessimism
- Punitiveness
- Self-Sacrifice
- Social Isolation
- Subjugation
- Unrelenting Standards
- Vulnerability

SESSION THREE: MODIFYING THOUGHTS AND BEHAVIORS

Self-Affirmations

This cheat sheet should help you create self-affirmations that are effective and personalized. Affirmations can focus on several categories, including character traits, physical attributes, and skills and/or accomplishments.

- 1. Start your affirmations with "I am" (e.g., "I am a good friend")
- 2. Keep them short (we don't need a novel)
- 3. Keep them positive (avoid saying "not")
- 4. Use feeling words when you can (e.g., "I am proud of myself")
- 5. Keep them focused on you (after all, they are **self**-affirmations)

Examples: "I am proud that I am hard-working" or "I am thankful for my strong legs"

My Self-Affirmations:

Feeling stuck? Ask yourself questions like these:

- When was a time that you felt proud of yourself and why?
- Have you ever forgiven someone who has hurt you?
- Have you ever received a compliment that you agree with?
- What would your best friend or a loved one say about you?

Feeling Guilty? Consider this:

- We are socialized to ignore or minimize our positive characteristics.
- Unfortunately, we then just dwell on our "negatives."
- Practicing self-affirmations helps bring balance and improve mood.

Alternative Response Worksheet

about?)	?? What? With whom? Wha e nt due for my class. I ask leclined.	
Alternative Thoughts and Images: (Are these thoughts helpful? Are the depressed thoughts 100% true/accurate, 100% of the time? What are other ways of looking at this? What is the bigger picture?)	Alternative Behaviors: (What could you do that would be more helpful for you, others, &/or the situation? What are coping strategies that might be helpful?) Coping Strategies I	Alternative Feelings: (What are feelings that are more helpful? What if you acted and thought differently about the situation? How might these changes help you feel differently?)
She might already have a partner. I've done well in this class so far, so I might not be stupid. Not everyone is going to like me and that is okay.	 Can Use: Deep breathing Distract myself Seek support from a friend/ family member Do a pleasurable activity Use alternative response worksheet Other Other 	Optimistic Confident Relaxed Neutral

Outcome: (What was the original outcome? Using these new alternatives, what would you like the outcome to be in the future?)

Original outcome: I left class early and e-mailed my professor to assign me a partner. Next time: I would like to say, "thanks anyway" and will ask a person with whom I have worked well in the past.

Alternative Response Worksheet

Situation (When? Where? What? With whom? What did you feel depressed about?)							
Alternative Thoughts and Images: (Are these thoughts helpful? Are the depressed thoughts 100% true/accurate, 100% of the time? What are other ways of looking at this? What is the bigger picture?)	Alternative Behaviors: (What could you do that would be more helpful for you, others, &/or the situation? What are coping strategies that might be helpful?) Coping Strategies I Can Use: Deep breathing Distract myself Seek support from a friend/ family member Do a pleasurable activity Use alternative response worksheet Other	Alternative Feelings: (What are feelings that are more helpful? What if you acted and thought differently about the situation? How might these changes help you feel differently?)					

Outcome: (What was the original outcome? Using these new alternatives, what would you like the outcome to be in the future?)



Challenging the Unhelpful Thinking Styles

If you're having trouble, ask yourself these questions:

Alternative THOUGHTS:

- 1. What are other ways of looking at this situation?
- 2. Am I looking at the whole picture?
- 3. What might be a more helpful way of thinking about this situation?
- 4. What unhelpful thinking styles might I be using here?
- 5. What is the evidence that my thoughts are true? Is there an alternative way of thinking about this situation that is more accurate?
- 6. What is the probability that my thoughts will happen? What are some other things that could happen that are equally, if not more, probable?
- 7. Have I had any experiences in the past that suggests that this thought might not be COMPLETELY true ALL of the time?
- 8. Can I really predict the future? Is it helpful to try? What is more helpful?
- 9. Am I exaggerating how bad the result might be? What is more realistic?
- 10. Can I read people's minds? What else might they be thinking?
- 11. If a friend or loved one were in this situation and had this thought, what would I tell them?


Alternative BEHAVIORS:

- 1. What could I do in the moment that would be more helpful?
- 2. What's the best thing to do (for me, for others, or for the situation)?
- 3. If my feared situation happens, how will I cope? What coping skills can I use to handle my feared situation? What have I done in the past that was successful?
- 4. Do I need to work on acceptance, letting go of control, being okay with less than perfect, or having faith in the future and myself?
- 5. Breathe: Focus your attention on your breathing. Imagine you have a balloon in your belly, inflating on the in-breath, deflating on the outbreath.

Alternative FEELINGS:

- 1. What might it feel like if I acted/thought differently?
- 2. When I'm not feeling this way, do I think about this situation differently?
- 3. Are there any strengths or positives in me or the situation that I might be ignoring?
- 4. What else might this feeling be related to? Is it *really* about feeling ____?
- 5. Tell yourself: "This feeling will pass. It's a normal body reaction."

Feelings Wheel



Other Helpful Thought Tips

1. **Stopping.** Interrupt a thought as it begins. Use a strong

image or a word to interrupt the thought. Strongly state it. either



internally or aloud.

- 2. Distracting. Redirect your mind to something else internally or externally, preferably something pleasant and engaging.
- 3. Mindful Observing. Watch,



label, or log your thoughts. Use the language "I am thinking ..." or "My mind is having the thought that..." to distance yourself from the thought.

- 4. **Understanding**. Begin to understand where thoughts come from by asking a number of questions, including: What is the purpose of this thought? Does it tie to a specific schema?
- 5. Mindfully Letting Go. Use imagery or words to visualize thoughts passing by. Good examples are clouds in the sky or leaves on a stream.
- 6. Gratitude. Try to focus on something that you are grateful for from the past, present, or future.

Helpful Behavioral Tips

The following lists of ideas are meant to get you thinking about possibilities for helpful behaviors. The activities do not have to be huge commitments; focus on small baby steps. Choose activities that are do-able for you when you are feeling down. The more success you have in doing the little things, the more likely you will be motivated to try more things.

Ideas for Socializing

- □ Watch a movie with a friend
- □ Go to an intermural or CP game
- □ Go to the library, UU, or dorm lounge to study

Ideas for Pleasant Activities

- □ Play with a pet
- □ Go shopping or window shopping
- □ Fix/tinker with something
- □ Listen to music

Ideas for Mastery Activities

- □ Wash a dish or two (even if they aren't yours)
- □ Do laundry
- □ Return a phone call
- \Box Write that email to your professor

- □ Go to a gym class, dance class, martial arts class, etc.
- □ Go eat free samples at Farmer's Market
- \Box Plan to eat a meal with a roommate/friend

- □ Color/Paint/draw/sculpt
- □ Write poem, music, play, story
- □ Read for fun
- □ Watch the sunset/rise at the beach

- \Box Read a chapter or a page of your homework assignment
- □ Take out the trash
- □ Pay a bill

Other Helpful Behavioral Tips for Managing Depression

1. **Get sunlight**: Aim for 5 to 15 minutes of sunlight a day.



Sunlight increases the brain's release of serotonin, which is associated with boosting mood.

- 2. **Get a massage**: Massage boosts serotonin and decreases stress hormones. Try giving yourself a massage by lying on or leaning against a tennis ball or rolling it against your muscles.
- 3. Practice Yoga: Yoga works to



improve mood and has benefits similar to that of exercise and relaxation techniques.

Yoga poses that incorporate back bends and opening the chest help to increase positive emotions.

- 4. **Stand up straight:** Your body tells your brain how to feel. Stand up straight and open your chest to feel more confident.
- 5. **Smile** (*even if you're faking*): When you smile, you are more likely to

perceive positive emotions in other people, which can positively impact your mood.



Smiling will lead your brain to believe that you are happy.

6. **Laugh** (*even if you're faking*): Your brain does not distinguish



between real and fake laughter. If your brain gets signals from your body that you should feel happy,

then it is more likely to feel happy.

- Do Progressive Muscle
 Relaxation: Flex a tight muscle for a few seconds and release. Pay particular attention to your facial muscles as those have the largest effect on emotion.
- 8. **Create/listen to music**: Music can help regulate your emotions. It can be soothing and help you feel calm and/or it can be energizing and pump you up.
- 9. **Dance**: Dancing combines music and physical activity so it is a double whammy.



10. **Journal:** Labeling your emotions and writing your story can be cathartic and help you organize your thoughts.

Adapted from Alex Korb's "The Upward Spiral"

My Personalized Plan

- 1. My schemas [located on pages **5-6**]:
- My depressive symptoms (e.g., lack of motivation, reduced sex drive) [located on page 7]:
- 3. My unhelpful thinking styles (e.g., catastrophizing) [located page 20]:
- 4. My unhelpful behaviors (e.g., isolating) [located on pages 18, 21]:
- 5. One sleep and one exercise tool I plan to use [located on **page 11**]:
- 6. Two thought exercises I plan to use (e.g., alternative responses, self-affirmations) [located on pages 31, 35, 37]:
- 7. Two behavioral tips I plan to use (e.g., pleasant event, mastery activity) [located on **pages 38-39**]:

APPENDIX

Understanding Depressive and other Related Disorders

While some depressive symptoms (i.e., sadness) are a normal experience for everyone, depressive and other related disorders are characterized by significant distress or impairment in social, academic/occupational, or other important areas of functioning (e.g., your general ability to function in life).

Some of the most common disorders include:

Major Depressive Disorder:	Involves experiencing depressive episodes, which must last for two weeks, and include a minimum of 5 symptoms.
Persistent Depressive Disorder:	A less extreme depressive presentation that includes a depressed mood and at least 2 other symptoms that last for at least two years. May have episodes of major depression along with periods of less severe symptoms, but symptoms must last for two years.
Premenstrual Dysphoric Disorder:	Several depressive symptoms present consistently during the week prior to menses.
Bipolar II Disorder:	Includes experiences of both depressive episodes as well as "mixed" or hypomanic episodes (i.e. elevated mood with other expansive symptoms).
Cyclothymic Disorder:	Persistent periods of alternating between "highs" and "lows" that are subthreshold for hypomanic and depressive episodes.
Disruptive Mood Dysregulation Disorder:	Recurrent temper outbursts and persistently irritable or angry mood for at least one year (present between ages 6-18).

Failure	Enmeshment/ Undeveloped Self	Vulnerability to Harm or Illness	Dependence/ Incompetence	Social Isolation/ Alienation	Defectiveness/ Shame	Emotional Deprivation	Mistrust/Abuse	Abandonment/ Instability	Schema
Does tasks in a halfhearted or haphazard manner	Lives through partner	Obsessively reads about catastrophes and anticipates them in everyday situations	Asks significant others (parents, spouse) to make all their financial decisions	Focuses exclusively on differences from others rather than similarities	Selects critical and rejecting friends; puts self down	Selects emotionally depriving partners and does not ask them to meet needs	Selects abusive partners	Selects partners who cannot make a commitment and remains in the relationships	Example of surrender
Avoids work challenges completely; procrastinates on tasks	Avoids intimacy; stays independent	Avoids going places that do not seem totally "safe"	Avoids taking on new challenges, such as learning to drive	Avoids social situations and groups	Avoids expressing true thoughts and feelings and letting others get close	Avoids intimate relationships all together	Avoids becoming vulnerable and trusting anyone; keeps secrets	Avoids intimate relationships; drinks a lot when alone	Example of avoidance
Becomes an "overachiever" by ceaselessly driving themself	Tries to become the opposite of significant others in all ways	Acts recklessly, without regard to danger	Becomes so self- reliant that they do not ask anyone for anything	Becomes a chameleon to fit into groups	Criticizes and rejects others while seeming to be perfect.	Acts emotionally demanding with partners and close friends	Uses and abuses others ("get others before they get you")	Clings to partner to point of pushing partner away; attacks partner for minor separations	Example of Over- Compensatio n
Punitiveness	Unrelenting Standards/ Hypercriticaln ess	Emotional Inhibition	Negativity/ Pessimism	Approval/ Recognition Seeking	Self-Sacrifice	Subjugation	Insufficient Self- Control/ Discipline	Entitlement/ Grandiosity	Schema
Treats self and others in harsh , punitive manner manner	Spends inordinate amounts of time trying to be perfect	Maintains a calm, emotionally flat demeanor	Focuses on the negative; ignores the positive; worries constantly;	Acts to impress others	Gives a lot to others and asks for nothing in return	Let's other individuals control situations and make choices	Gives up easily on routine tasks	Bullies others into getting own way, brags about own accomplishments	Example of Surrender
Avoids others for fear of punishment	Avoids or procrastinates in situations where performance will be judged	Avoids situations in which people discuss or express feelings	Drinks to blot out pessimistic feelings and unhappiness	Avoids interacting with those whose approval is coveted	Avoids situations involving giving or taking	Avoids situations that might involve conflict with another individual	Avoids employment or accepting responsibility	Avoids situations in which they are average	Example of Avoidance
Behaves in overly forgiving way	Does not care about standards at all-does tasks in a hasty, careless manner	Awkwardly tries to be the "life of the party, "even though it feels forced and unnatural	ls overly optimistic denies unpleasant realities	Goes out of the way to provoke the disapproval of others; stays in the background	Gives as little to others as possible	Rebels against authority	Becomes overly self- controlled or self- disciplined	Attends excessively to the needs of others	Example of Over- Compensation