## A benefit of high temperature: increased effectiveness of a rice bacterial blight disease resistance gene

## K. M. Webb<sup>1,2</sup>, I. Oña<sup>2</sup>, J. Bai<sup>1</sup>, K. A. Garrett<sup>1</sup>, T. Mew<sup>2</sup>, C. M. Vera Cruz<sup>2</sup> and J. E. Leach<sup>3</sup>

<sup>1</sup>Department of Plant Pathology, Kansas State University, 4024 Throckmorton Plant Sciences, Manhattan, KS 66506, USA; <sup>2</sup>Plant Breeding, Genetics and Biotechnology, The International Rice Research Institute, Los Baños, Philippines; <sup>3</sup>Department of Bioagricultural Sciences and Pest Management, Plant Sciences, Colorado State University, Ft Collins, CO 80537-1177, USA

## Summary

Author for correspondence: Jan E. Leach Tel: +1 (970) 491 2924 Email: jan.leach@colostate.edu

Received: 29 May 2009 Accepted: 19 September 2009

*New Phytologist* (2010) **185**: 568–576 **doi**: 10.1111/j.1469-8137.2009.03076.x

Key words: aggressiveness, avirulence/effector gene, durability, fluctuating selection, *Oryza sativa* (rice), oscillating selection, virulence, *Xanthomonas oryzae* pv. *oryzae* (Xoo). • Continuous planting of crops containing single disease resistance (R) genes imposes a strong selection for virulence in pathogen populations, often rendering the R gene ineffective. Increasing environmental temperatures may complicate R-gene-mediated disease control because high temperatures often promote disease development and reduce R gene effectiveness. Here, performance of one rice bacterial blight disease R gene was assessed in field and growth chamber studies to determine the influence of temperature on R gene effectiveness and durability.

• Disease severity and virulence of *Xanthomonas oryzae* pv. *oryzae* (*Xoo*) populations were monitored in field plots planted to rice with and without the bacterial blight *R* gene *Xa7* over 11 yr. The performance of *Xa7* was determined in high-and low-temperature regimes in growth chambers.

• Rice with Xa7 exhibited less disease than lines without Xa7 over 11 yr, even though virulence of Xoo field populations increased. Xa7 restricted disease more effectively at high than at low temperatures. Other R genes were less effective at high temperatures.

• We propose that Xa7 restricts disease and Xoo population size more efficiently in high temperature cropping seasons compared with cool seasons creating fluctuating selection, thereby positively impacting durability of Xa7.

## Introduction

Climate change presents considerable challenges for plant disease management (Coakley et al., 1999; Garrett et al., 2006). While the specific effects of climate change on plant diseases will depend on the plant-pathogen system in question, the impact of some changes, such as temperature and precipitation, can be predicted for several plant species from empirical studies (reviewed by Coakley et al., 1999). For example, increased temperature results in higher susceptibility of wheat and oats to rust diseases, while forage species become more resistant to certain fungi (Coakley et al., 1999). The impact of increased temperatures associated with climate change on plant disease are speculated to be through both the host plant and the pathogen. These proposed mechanisms for effects, as reviewed by Coakley et al. (1999) and Garrett et al. (2006), include alterations of plant architecture to create more suitable microclimates for pathogen colonization and disease development; changes in

the function and effectiveness (ability to restrict pathogen multiplication and colonization) of plant disease resistance; and increased abiotic stresses on plants, which would affect susceptibility to plant pathogens.

Higher temperatures are predicted to accelerate the breakdown of plant disease resistance through higher disease pressure and/or altered resistance gene efficacy in many host-pathogen systems. Higher disease pressure is coincident with elevated pathogen populations, and this, in turn, increases the probability that a mutation will allow the pathogen population to overcome a single resistance (R) gene, the primary sources of resistance in many crop breeding programs. There are examples of R genes that are either more or less effective at high temperatures (Dyck & Johnson, 1983; Browder & Eversmeyer, 1986; Kolmer, 1996; Wang *et al.*, 2001; Eizenberg *et al.*, 2003; Uauy *et al.*, 2005). In the *Puccinia recondite*—wheat pathosystem, some, but not all, wheat resistance genes are more effective at higher temperatures, and this temperature effect is highly

isolate-dependent (Dyck & Johnson, 1983; Browder & Eversmeyer, 1986; Kolmer, 1996) The wheat stripe rust R gene *Yr36* confers resistance to a broad spectrum of races of the pathogen *P. striiformis* f. sp. *tritici* at high temperatures (25–35°C), while at low temperatures (15°C), wheat with *Yr36* is susceptible to the fungus (Uauy *et al.*, 2005). The recent cloning of *Yr36* may facilitate understanding of how the gene function is influenced by temperature (Fu *et al.*, 2009).

Deciphering the interconnectivity of temperature, pathogen population evolution and R gene effectiveness will require an understanding of how *R* genes function and how that function is affected by high temperature. In general, Rgenes directly or indirectly interact with the products of pathogen effector genes (also called avirulence or *avr* genes), and this interaction signals a disease resistance cascade (for review, see Chisholm et al., 2006). Effector proteins function as critical components of a pathogen's virulence, and thus plants have evolved to recognize the effectors as a trigger for defense. This suggests that a mutation in an effector gene can result in a loss of R gene recognition and increased host range, but their importance to virulence means that mutation can also result in reduced pathogenic fitness (Leonard & Czochor, 1980; Leach et al., 2001; Burdon & Thrall, 2003). Pathogenic fitness, in this context, refers to aggressiveness to a susceptible host, and ability to persist in the population. Thus, the more important an effector gene is to a pathogen's fitness, the more difficult it will be for the pathogen to adapt to overcome the corresponding plant Rgene (Leach et al., 2001). While there are field and laboratory studies that support the prediction that R gene durability correlates with the importance of pathogen effectors to a pathogen's fitness (reviewed in Leach et al., 2001), there are no studies that integrate the influence of climate, particularly high temperatures, with pathogen population evolution and *R* gene function or durability.

Understanding the interaction of temperature and *R* gene function and durability is complicated in systems with multiple crops per year, such as rice, where temperatures can vary between being conducive and nonconducive for disease by season. Standard definitions of resistance gene durability include the continuing effectiveness of the gene when environmental conditions are conducive to disease for long periods of time (Johnson, 1984), but seasonal breaks in disease conduciveness can make the useful life of resistance genes longer in practice. Changing environmental conditions can impose a fluctuating selection pressure on pathogen populations (Lynch, 1987; Hairston & Dillon, 1990). Indeed, seasonality may produce regular fluctuations in environmental conditions, or oscillations, allowing for a reversal in the direction of selection (Gibbs & Grant, 1987; Grant & Grant, 2002). While not yet investigated, the influence of such fluctuating selection on R gene durability will likely vary with *R* gene, and will depend on the speed of pathogen evolution to races that can overcome resistance, as influenced by changes in infection cycles and the availability of suitable microclimates for disease development.

We have been studying the factors influencing the durability of plant R genes in rice to the bacterial blight (BB) pathogen, Xanthomonas oryzae pv. oryzae (Xoo). In a previous field study, the cost of pathogenic fitness for Xoo was shown to be an important factor contributing to the durability of one rice BB R gene, Xa7 (Vera Cruz et al., 2000). The effectiveness of the Xa7 gene was correlated with mutations in Xoo that resulted in reduced pathogenic fitness, as measured by reduced aggressiveness on susceptible rice cultivars (Bai et al., 2000; Vera Cruz et al., 2000). These mutations occurred specifically in the pathogen effector gene avrXa7, which is important to pathogenic fitness (Bai et al., 2000; Ponciano et al., 2004). By contrast, mutation of the effector gene avrXa10 did not affect pathogenic fitness, and the corresponding R gene Xa10 was rapidly overcome in the field (Bai et al., 2000; Vera Cruz et al., 2000). Thus, selection on the pathogen population imposed by the Xa7 gene, but not the Xa10 gene, resulted in reduced pathogenic fitness (Bai et al., 2000; Ponciano et al., 2004), and the cost of pathogenic fitness to the pathogen was one factor in effectiveness of the Xa7 gene (Vera Cruz et al., 2000).

Although the costs of pathogenic fitness affected pathogen population structure when under R-gene-imposed selection, other factors are likely to also influence an R gene's durability. In the BB-rice system, temperature is a likely influence because there are typically two cropping seasons in tropical regions, a cool season and a hot season (Peng et al., 2004). At the International Rice Research Institute (IRRI) farms in the Philippines, for example, between 1979 and 2003, cool-season temperatures averaged a maximum of 30°C and a minimum of 22.4°C (extreme temperatures of 31.4 and 21.0°C), and the hot-season temperatures averaged a maximum of 31.3°C and a minimum of 24.4°C (extreme temperatures of 35.7 and 16.4°C) (Centeno et al., 1995; Peng et al., 2004). High temperature is conducive to BB development in field situations, and disease severity is particularly high in the hot season (Yamada et al., 1979; Horino et al., 1982; Ou, 1985; Ezuka & Kaku, 2000). By contrast, little disease is observed in the cool season.

In our previous experiments, after 3 yr continuous planting with Xa7 in the same field sites (six cropping seasons), we measured an increase in a Xoo population that could cause disease on rice with Xa7 (loss of R gene recognition) without loss of pathogenic fitness (aggressiveness function). This group of strains, designated race 9b, exhibited a wide variation in their aggressiveness to rice, including rice with Xa7 (Ponciano *et al.*, 2004). The detection of the race 9b strains in the fields raised the possibility that the Xoo population was evolving to one that can overcome resistance controlled by Xa7 while maintaining fitness on susceptible hosts.

In this study, we monitored the structure of the field Xoo population for an additional 9 yr (total 11 yr) to determine if the Xoo strains that had lost avrXa7 function but maintained pathogenic fitness would break down resistance conferred by Xa7. We show that, although the Xoo population virulent to Xa7 dominated the field population, Xa7-directed resistance remained effective. The differences between disease pressure observed in the hot (conducive) and cool (nonconductive) rice cropping seasons and the observations from other host-pathogen systems that some R genes differ in their effectiveness at high temperatures led us to hypothesize that Xa7 may be more effective at high temperatures, and that this is one factor contributing to the relative durability of the R gene. As a first step to testing this hypothesis, we demonstrate in this study that Xa7 is more effective at high than at low temperatures relative to other BB R genes. We speculate that the greater effectiveness of Xa7 in the hot season may slow the evolution to more aggressive races of Xoo in the field. A novel aspect of this work is that, because field plots were established and maintained at one epidemiological location with continuous rice crops during both growing seasons in the Philippines from 1993 to 2004, long-term analysis of the impacts of the Xa7 gene on evolution of X00 populations under fluctuating selection pressures imposed by temperature was possible.

#### Materials and Methods

#### Field experimental design and disease assessment

Experiments were conducted in neighboring field plots located in Calauan, the Philippines, during the years 1993-95, 1998-99 and 2002-04 as described previously (Vera Cruz et al., 2000). During both the cool and hot seasons, rice lines were planted in a randomized complete block design using  $13 \times 13$  m plots with three (1998–99) or four (1993-95 and 2002-04) replications. In 1993-95, rice with the Xa7 R gene (IRBB7) was present in 20% of the plots, and the remaining 80% of the plots were planted with nearisogenic rice lines that did not contain Xa7, including the recurrent susceptible parent, IR24. In 1998-99, 33% of the plots contained Xa7, and in 2002-04, 60% of the plots contained Xa7. Data on the monthly average temperatures were collected from 1993 to 2004 (Sheehy et al., 2005). Disease estimates and leaf samples were collected in a W pattern throughout each plot, with either seven (1993–95) or five (1998-2004) collection sites per W pattern (Vera Cruz et al., 2000). Disease estimates and leaf samples were taken from IRBB7 and IR24 (lacks Xa7) plots in the hot season only, as little disease is observed during the cool season (Ou, 1985; Ezuka & Kaku, 2000). Estimates of disease, including incidence (occurrence of symptoms) and disease severity (% diseased leaf area, % DLA), were assessed and bacteria were isolated from sampled symptomatic tissue as

previously described (Oña *et al.*, 1998; Vera Cruz *et al.*, 2000). Differences in disease incidence and severity were evaluated in an analysis of variance (ANOVA) using Proc Mixed in SAS statistical software (SAS Institute, version 9.2, Cary, NC, USA).

#### Xoo population structure

Changes to X00 populations in plots planted to rice line IR24, which does not contain Xa7, were monitored by assessing the race structure and aggressiveness patterns for strains collected from these plots over three time frames (1993-95, 1998-99, and 2002-04). Up to 15 bacterial strains per plot per replication per year were randomly selected and the race of each strain was determined by inoculation to a set of seven rice differential lines, each containing a different BB R gene or no known R genes (IR24, susceptible control) (Kauffman et al., 1973; Ogawa et al., 1991). Changes in the Xoo population structure in terms of both aggressiveness (to susceptible host) and virulence (to resistant host) over time were measured by calculating the mean lesion lengths for all strains collected within a single year after inoculation to IR24 and IRBB7, respectively. Xoo race 3 and races 9b and 9a/c are differentiated by lesion lengths on IRBB7 as follows: race 3, 0-4.9 cm; race 9b, 5-17 cm; and race 9a/c, > 17 cm. To identify differences in X00 population aggressiveness and virulence, the mean lesion lengths were evaluated using ANOVA in Proc Mixed.

#### Effect of temperature on R gene function

To determine the effect of temperature on various R genes, near-isogenic rice lines IRBB3, IRBB4, IRBB5, IRBB7, and IRBB10, which contain Xa3, Xa4, xa5, Xa7 and Xa10, respectively, as well as the recurrent susceptible parent IR24 (Ogawa et al., 1991), were grown in a glasshouse for 11 wk. Near-isogenic lines were used to minimize differences owing to genetic background and so that all effects could be associated directly with the R gene in question. One week before inoculation, the plants were transferred to growth chambers with 70% RH and day : night temperatures of 35:27 and 29:21°C, respectively. Xoo strains (races) used for inoculation induced resistance in combination with each R gene, and were PXO61 (race 1), PXO79 (race 3), PXO71 (race 4), and PXO112 (race 5) (Mew, 1987). Flag leaves were inoculated with bacteria (concentration was adjusted to c.  $5 \times 10^8$  CFU ml<sup>-1</sup>) at 85 d after sowing (das) using a pin-prick method as described in (Horino et al., 1982; Ou, 1985). Lesion lengths were measured 21 d after inoculation (dai). Experiments were performed using a split-plot design with temperature applied at the whole-plot level and variety-isolate combinations applied at the subplot level (three replications, two plants

per replicate). The experiment was performed three times and differences in lesion length from isolates causing resistant interactions for each R gene were assessed using SAS Proc Mixed. In a separate experiment, IRBB7 and IR24 plants grown at 35 : 29 and 29 : 21°C were inoculated using a leaf clip assay (Kauffman *et al.*, 1973) with two race 3 isolates (PXO340 and PXO368). Plants were inoculated at 45 das, and lesion lengths measured at 14 dai. Lesion lengths from both isolates on a single line were combined and analyzed for significant differences using ANOVA (SAS Proc Mixed).

To comprehensively evaluate the effects of different temperature regimes (35:31 and 29:21°C) on interactions with Xa7, strains of race 9a/c without avrXa7 (PXO346, PXO2684), race 3 with avrXa7<sup>+</sup> (PXO1865, PXO368), and race 9b with partially functional avrXa7 (PXO348, PXO0314, PXO350, PXO354, PXO355, PXO356, PXO360 and PXO554) were used to inoculate IRBB7 and IR24 at 45 das as previously described (Kauffman et al., 1973). A split-plot design was used, with temperature applied at the whole-plot level, and variety-isolate applied at the subplot level (two replicates, with two plants per replicate). The experiment was performed twice. Lesion lengths were measured at 3, 6, 9, 12, 15, and 18 dai. Area under the disease progress curves (AUDPC) were calculated (Madden et al., 2007) and significant differences compared using ANOVA (SAS Proc Mixed).

## Results

Evolution of *Xoo* populations on a susceptible host over 11 yr

To estimate the impact of deploying rice with Xa7 over 11 yr on the race structure of Xoo field populations, the race type of Xoo isolates from the susceptible host (IR24) was determined. IR24 acts as a susceptible 'trap' plot as it will produce disease with all indigenous Xoo races that are present in the local population, and the Xoo population on IR24 reproduces in the absence of selection from any Rgene being deployed in other plots. In 1993 and 1994, Xoo strains in the field plots were predominantly race 3 (Fig. 1). Race 2 strains, which dominated in the population in the Philippines before 1993 (Mew et al., 1992; Vera Cruz et al., 1996), were only detectable in early samples, and were not detected on IR24 after 1999. Race 9 strains were distinguished into two functional subgroups (race 9b, and 9a/c) based on the function of the Xoo avrXa7 gene (Mew et al., 1992; Vera Cruz et al., 2000). Both subgroups were isolated from the susceptible host IR24, with race 9b populations significantly increasing (starting in 1995), and then stabilizing as the dominant component of the population (Fig. 1). Concurrently, the prevalence of race 3 (recognized by Xa7) decreased. Significant increases in the proportion of





**Fig. 1** Frequency (% of total population) of *Xanthomonas oryzae* pv. *oryzae* races (2, 3, 9b, 9a/c) in the population isolated from rice with no *R* gene (IR24) in field sites located in Calauan, the Philippines, over 11 yr (22 cropping seasons). Lower-case letters indicate significant differences in race frequency within each year with separations calculated based on an *F*-test in SAS Proc Mixed (P = 0.05).

race 9a/c (22.5%; not recognized and therefore virulent to Xa7 as a result of mutation of avrXa7 function) were detected starting in 1998 (Fig. 1). By 2004, race 9, including both subgroups (9b and 9a/c), was the dominant race in the field, with the incidence of subgroup 9a/c increasing to 31.1% of the population. Thus, from 1993 to 2004, the *Xoo* population structure at these field sites changed significantly from dominance by strains that are controlled by Xa7 (race 3) to a population dominated by strains able to cause disease (virulent) on rice with Xa7 (race 9), as indicated in a mean separation test (least significance difference) of the yearly means (P = 0.05).

# Virulence and aggressiveness increased in the *Xoo* population

In previous studies, a loss of Xoo avrXa7 gene function resulted in reduced aggressiveness (less disease) to susceptible rice hosts, suggesting reduced pathogenic fitness (Bai et al., 2000; Vera Cruz et al., 2000). Because the Xoo field population changed to being primarily composed of strains able to cause disease on rice with Xa7 (race 9) (Fig. 1), we asked if the overall population was more or less aggressive to IR24. Xoo strains were collected from each year of the study, and aggressiveness was determined by measuring the disease lesion length on IR24 and IRBB7 (Fig. 2). Lesion lengths provide a measure of pathogen colonization of the rice tissues (Barton-Willis et al., 1989). In 1993, the mean



**Fig. 2** The Xanthomonas oryzae pv. oryzae population increases in virulence to rice with Xa7 (IRBB7; rectangles) and maintains aggressiveness to IR24 (triangles) over three time periods from 1993 to 2004. Mean comparisons were done for each period (underlined) and year within a period. Upper-case letters are for comparisons with IR24, while lower-case letters are for IRBB7 and indicate population means that are significantly different based on an *F*-test in SAS Proc Mixed (P = 0.05).

lesion lengths of the *Xoo* population on IRBB7 and IR24 were 4.6 and 20.5 cm, respectively, which is consistent with a functional *Xoo avrXa7* gene (Mew *et al.*, 1992) in the predominant race 3 population. Between 1995 and 2004, the mean lesion length of the *Xoo* population on rice with *Xa7* significantly increased from 10.5 to 15.1 cm. The population's aggressiveness to IR24 also increased between 1993 and 1995 and then remained stable through 2003, with a mean lesion length of *c.* 26 cm (Fig. 2). This increase in aggressiveness in the overall pathogen population is associated with a decrease in race 3 and an increase in race 9 in the population (Fig. 1).

#### Disease incidence and severity

To determine if the increase in virulent race 9 strains in the *Xoo* population correlated with an increase in disease on rice with *Xa7*, we assessed disease development in the rice plots over the 11 yr period. Disease incidence (% of plants showing lesions) and severity (% DLA) were monitored on rice with and without *Xa7* during the hot seasons of 1993–95, 1998–99, and 2002–04 (Table 1). Although both disease incidence (48–94%) and severity (5–32%) were consistently high on rice without *Xa7*, both indicators of disease were significantly lower on IRBB7 than on the susceptible IR24, for all years included in the study (Table 1); this indicates that resistance conferred by *Xa7* was still effective during each year of the study. Even though the proportion of

Table 1Significant terms from an ANOVA of the disease severity(% diseased leaf area) and disease incidence (% occurrence ofsymptoms) of bacterial blight in field plots from the hot season(1993–2004)

Year	Disease severity		Disease incidence	
	IR24	IRBB7	IR24	IRBB7
1993	32.04a	0.68b	94.44a	15.76b
1994	21.12a	0.35b	89.74a	14.93b
1995	5.68a	0.15b	66.67a	7.64b
1998	26.37a	1.68b	50a	3.33b
1999	18.33a	2ba	48.33a	8.33b
2002	4.5a	0.43b	84.5a	44.1b
2003	6.88a	2.63b	73.65a	46.15a*
2004	5.2a	0.57a	82.23a	19.27b

Values are means of four replications (1993–95; 2002–03) or three replications (1998–99, 2004). Means within a row by year followed by a different letter are significantly different ( $\alpha = 0.05$ ) based on an *F*-test in SAS Proc Mixed.

\*Not significantly different ( $\alpha = 0.05$ ), P = 0.0579.

the pathogen population virulent to Xa7 (race 9) was as high as 100% in 2004 (Fig. 1), BB incidence and severity remained low for IRBB7 (Table 1). Thus, although race 9 strains became the predominant population, Xa7 continued to effectively restrict BB disease in the field. These results also suggested that additional factors outside pathogenic fitness contributed to the continued effectiveness of Xa7.

## *Xa7*-mediated resistance is more effective at high temperatures

Although race 9 strains (virulent to Xa7) are predominant in the field plots, rice varieties with Xa7 continued to exhibit less disease than IR24 (Table 1). During the initial analysis (in 1998) of Xoo population race structure, we tested field strains on IR24 and IRBB7 in March/April (average temperature 33°C) and January (average temperature 29°C) under screenhouse conditions. A two-tailed *t*-test, assuming unequal variances, showed that the field strains caused shorter lesions on IRBB7 when screened in months with higher temperatures than in months with cooler temperatures ( $P = 9.0 \times 10^{-14}$ ) (Fig. 3). This suggested that the Xa7 gene was more effective at high temperatures than at low temperatures.

To confirm that Xa7 is more effective at high temperature and to determine if the effect is unique to Xa7, performances of five BB *R* genes in high- (35°C day : 27°C night) and low-temperature (29°C day : 21°C night) regimes were compared in growth chamber studies (Fig. 4a). Four of the five genes examined – Xa3, Xa4, xa5, and Xa7 – were affected by temperature in some way. Xa3, Xa4, and xa5 were less effective at high temperatures (as indicated by longer lesions) than at low temperatures (Fig. 4a). In particular, Xa4 exhibited much longer lesions at high



**Fig. 3** Mean lesion lengths for a population of *Xanthomonas oryzae* pv. *oryzae* strains that were inoculated to IRBB7 and IR24 at two different time periods (January 1998 (black squares) and March/April 1998 (gray diamonds)). Average daytime temperatures were 29 and 33°C, respectively. Lesions were generally shorter on IRBB7 during the higher temperatures (April). Population means were significantly different using a two-tailed *t*-test (P < 0.05).



**Fig. 4** *Xa7* is more effective in higher (gray bars) than in lower (black bars) temperature regimes. (a) Leaves of rice near-isogenic lines with *R* genes *Xa3*, *Xa4*, *xa5*, *Xa7*, and *Xa10* were inoculated with strains of *Xanthomonas oryzae* pv. *oryzae* that result in resistant interactions using a pin-prick method and grown in high-(35 : 27°C, day : night) and low-temperature regimes (29 : 21°C, day : night). (b) Leaves of rice near-isogenic lines containing the *Xa7* gene (IRBB7) and no *R* gene (cultivar IR24) were inoculated using a leaf clip method grown in high- (35 : 29°C, day : night) and low-temperature (29 : 21°C, day : night) regimes. Lower-case letters indicate significant differences between temperature regimes within a single *R* gene interaction (*P* = 0.05).

temperatures. By contrast, Xa7 was more effective (produced shorter lesions) at the high temperatures. The increased effectiveness of Xa7 at higher temperatures was confirmed using a second inoculation assay, the leaf clip inoculation method (Kauffman *et al.*, 1973) (Fig. 4b). In these experiments, there was no significant difference in the amount of disease on the near-isogenic susceptible host IR24.

The possible influence of bacterial genotype on increased function of Xa7 at high temperatures was tested by inoculating several Xoo strains with nonfunctional (race 9a/c), partially functional (race 9b), or functional (race 3) avrXa7 alleles to IRBB7 and IR24. We used the AUDPC as a measure of the rate of disease development that includes lesion length measurements over time (Madden et al., 2007). At high temperatures, the amount of disease on IRBB7 induced by all Xoo isolates with functional or partially functional avrXa7 alleles was reduced (average AUDPC, 74.2) relative to low temperatures (average AUDPC, 142.9) (Fig. 5a, upper case letters). Because the *avrXa7* allele varies among these bacterial strains (Bai et al., 2000; Ponciano et al., 2004), and this can contribute to differences in AU-DPC on Xa7, we tested if there was an effect of temperature on the avrXa7 gene in particular strains. In this study, we did not directly partition the effect of temperature on the pathogen *avr* effector gene vs the rice R gene; however, there was no temperature  $\times$  strain  $\times$  variety interaction (P = 0.857), nor a temperature  $\times$  strain interaction (*P* = 0.899), suggesting that temperature does not have an impact on the function of the pathogen avrXa7 gene. Additionally, there was no difference in AUDPC ( $\alpha = 0.05$ ) when each strain was compared with itself at either temperature or in each of the varieties tested (data not shown). The two race 9a/c strains (no avrXa7 function) (Ponciano et al., 2004) cause about the same amount of disease at both temperatures (Fig. 5a,b). Race 3 and race 9b strains cause less disease than race 9a/c strains on IRBB7 independent of temperature, and at high temperature there was no difference in AUDPC caused by most race 3 and 9b strains. These data confirm that Xa7 is more effective at higher temperatures. Furthermore, the data suggest that, at the high temperature, the R gene more efficiently recognizes pathogen strains in which the *avrXa7* alleles either are partially mutated or have complete function, as these strains cause intermediate responses at the low temperature (as compared with isolates that have completely lost avrXa7 function at low temperatures) (Fig. 5a).

An ANOVA did not reveal differences in AUDPC at the two temperature regimes on the susceptible host (IR24, Fig. 5b). However, a sign test was performed to determine if the mean AUDPCs for the low- and high-temperature regimes differed, and it indicated that, when each of the 13 isolates was evaluated as a separate trial, significantly longer lesions were produced at higher temperatures than were produced at lower temperatures (P = 0.0001) on susceptible hosts, supporting the previous studies (Ou, 1985; Ezuka & Kaku, 2000).



**Fig. 5** Xa7 is more effective in higher (35 : 31°C, day : night) than in lower (29 : 21°C) temperature regimes. Area under the disease progress curves (AUDPC) for rice cultivar IRBB7 (a) and cultivar IR24 (b) after inoculation with 13 strains of *Xanthomonas oryzae* pv. *oryzae*, representing race 3 (functional *avrXa7*; gray bars), race 9b (variation in *avrXa7* function; white bars) and race 9a/c (nonfunctional *avrXa7*; black bars). Inoculation was by the leaf clip method. Lower-case letters indicate significant differences (P < 0.05) among the strains within a single temperature regime. Upper-case letters indicate significant differences among all the strains comparing the two different temperature regimes within a cultivar.

#### Discussion

One factor influencing the durability of the BB R gene Xa7 is the fitness penalty associated with the loss of the corresponding pathogen effector gene function (avrXa7) (Vera Cruz et al., 2000). This would imply that the Xoo population would slowly develop increased virulence to Xa7 only after extended deployment of the R gene. Over the 11 yr (1993-2004) of our field study, the population of Xoo isolated from the susceptible host (IR24) increased in virulence to rice with Xa7 (Fig. 1). This change was attributed to the increase in the proportion of Xoo races 9b and 9a/c, which have reduced or caused complete loss of avrXa7 function, respectively, and are virulent to rice with Xa7 (Ponciano et al., 2004; Fig. 1). The Xoo field population from 1995 onwards also caused longer lesions on rice with or without Xa7 relative to earlier years (Fig. 2). The increase in aggressiveness was attributed to the increase in predominantly race 9b Xoo strains that had partially lost avrXa7 function, but had not lost pathogenic fitness. What is most intriguing is that, despite the increased pathogen virulence to Xa7, disease severity on IRBB7 remained low during the 11 yr period and that resistance remained effective (Table 1). These results confirmed that pathogenic fitness cost is only one factor in pathogen evolution to virulence, and pointed to the need to explore other factors that might also influence Rgene durability.

An R gene is only considered to be durable after it has been effective while deployed over a long time, in several environments, and under conditions conducive to disease development (Johnson, 1984). In the Philippines, rice is grown in two seasons with different environmental growing conditions, a dry, cool season and a wet, hot season. Because environment is predicted to influence *R* gene durability, we tested the impact of one environmental factor, high temperature, on *Xa7* function. We found that *Xa7* functions more effectively at higher temperatures, that is, it restricts the development of lesions (shorter lesions) on leaves compared with lower temperatures (Figs 4,5). The average maximum temperature during the hot season at the field site from 1979 to 2003 was *c*. 31.1°C, with high temperature extremes of > 35°C (Peng *et al.*, 2004; Sheehy *et al.*, 2005). We predict that if only *Xa7* had been deployed in the plots, and the susceptible IR24 plants had not been there to serve as a reservoir for the race 9b populations, the greater effectiveness of *Xa7* at high temperatures would have reduced the race 9b populations much more over the 11 yr period.

We propose that Xoo field populations developing on Xa7 experience fluctuating selection pressure in the shift from the hot to cool seasons. During the hot season, the abiotic environment is more conducive to disease. The greater effect of Xa7 during the hot season can have a complicated effect on selection for virulence in Xoo: selection for virulent individuals within the population is stronger than during the cool season because of the greater effect of Xa7, but population size is reduced compared with the hot season potential. The pathogen numbers that would carry over into the cool season would be substantially reduced. Thus, we suggest that when the cool season crop is planted, the reduced initial inoculum load, the presence of Xa7 (even if reduced in effectiveness), and the cool, nonconducive temperatures would restrict disease development and pathogen numbers further. The effect of these two alternating forms of selection on the X00 population across years is likely to increase the effective life of Xa7. Unfortunately, we were

not able to measure the amount of inoculum at the beginning of the cool season, nor could we measure disease in that season, because little or no disease was observed in the cool season.

While the BB R gene Xa7 was more effective when incubated at high temperatures, the opposite was true for BB Rgenes Xa3, Xa4, and xa5. These differential responses of R genes to temperature may provide additional information when predicting R gene durability. In our study, Xa10 resistance was not affected by temperature. This, combined with the fact that the corresponding avrXa10 effector gene in Xoo was found to be unimportant for pathogenic fitness (Bai et al., 2000), may explain why Xa10 was not durable in the field (Vera Cruz et al., 2000). Intriguingly, although Xa4 was much less effective at higher temperatures, it confers durable resistance in many countries (Mew, 1987; Bonman et al., 1992). This R gene does not exhibit a classic hypersensitive response (Guo et al., 1993) and is predicted to exhibit a quantitative rather than a qualitative type resistance (Li et al., 2001), which may influence its durability. Because it has not yet been cloned, the contribution of the avrXa4 gene to pathogenic fitness, if any, is currently unknown.

The biological basis for enhanced activity of Xa7 at high temperatures is not known. The rice blast R gene Pib is upregulated at high temperatures, and it is predicted that the increased production of the R gene product enhances defense response (Wang *et al.*, 2001). Increased amounts of the Xa7 protein at high temperatures could account for enhanced recognition of the wild-type and mutant AvrXa7 effector proteins. Alternatively, protein–protein interactions involving Xa7 may be affected by high temperatures. These hypotheses cannot be tested currently as the Xa7 gene has not been cloned.

Previous work demonstrated that the fitness cost imposed on the pathogen to overcome resistance can influence R gene durability (Vera Cruz et al., 2000). We have shown that another likely factor influencing R gene durability is the enhanced effectiveness of an R gene at high temperature. While we have only tested the cost of pathogenic fitness (Vera Cruz et al., 2000) and the effect of temperature, it is likely that other factors, such as humidity and nutrition, as well as host genetic background and developmental attributes (Mazzola et al., 1994; Cao et al., 2007; Iyer-Pascuzzi et al., 2008), will influence R gene durability, and that not all factors will affect all R genes or have an equal effect on them (Burdon & Thrall, 2003). We predict that temperature affects R gene durability by causing variable (fluctuating) selection on the pathogen population between the cool and hot growing seasons. Given the myriad ways that climate change will affect plant disease management (Garrett et al., 2006), it is important to include both environmental factors, such as temperature, and pathogenic fitness parameters in the development of predictive models that will accurately identify durable *R* genes before their deployment.

## Acknowledgements

This research was supported by USDA-CSREES-NRI #2003-01551, a USAID Linkage Project, and the Kansas State and Colorado State Agricultural Experiment Stations. KMW was also supported by the Asia Rice Foundation and a James B. Person Fellowship. Field studies were supported by IRRI. We thank T. Loughlin and B. Bartolome for statistical assistance, and C. Jahn, T. Adhikari and I. Eujayl for critically reading the manuscript.

## References

- Bai JF, Choi SH, Ponciano G, Leung H, Leach JE. 2000. Xanthomonas oryzae pv. oryzae avirulence genes contribute differently and specifically to pathogen aggressiveness. *Molecular Plant-Microbe Interactions* 13: 1322–1329.
- Barton-Willis PA, Roberts PD, Guo A, Leach JE. 1989. Growth dynamics of Xanthomonas campestris pv. oryzae in leaves of rice differential cultivars. *Phytopathology* 79: 573–578.
- Bonman JM, Khush G, Nelson R. 1992. Breeding rice for resistance to pests. *Annual Review of Phytopathology* 30: 507–528.
- Browder LE, Eversmeyer ME. 1986. Interactions of temperature and time with some *Puccinia recondita: Triticum* corresponding gene pairs. *Phytopathology* 76: 1286–1288.
- Burdon JJ, Thrall PH. 2003. The fitness costs to plants of resistance to pathogens. *Genome Biology* 4: 227.
- Cao Y, Ding X, Cai M, Zhao J, Lin Y, Li X, Xu C, Wang S. 2007. The expression pattern of a rice disease resistance gene *xa3/xa26* is differentially regulated by the genetic backgrounds and developmental stages that influence its function. *Genetics* 177: 523–533.
- Centeno HGS, Balbarez AD, Fabellar FG, Kropff MJ, Matthews RB. 1995. Rice production in the Philippines under current and future climates. In: Matthews RB, Kropff MJ, Bachelet D, van Laar HH, eds. *Modeling the impact of climate change on rice production in Asia.* CAB International, Wallingford and the International Rice Research Institute, Manila, 237–250.
- Chisholm ST, Coaker G, Day B, Staskawicz BJ. 2006. Host-microbe interactions: shaping the evolution of the plant immune response. *Cell* 124: 803–814.
- Coakley SM, Scherm H, Chakraborty S. 1999. Climate change and disease management. *Annual Review of Phytopathology* 37: 399–426.
- Dyck P, Johnson R. 1983. Temperature sensitivity of genes for resistance in wheat to *Puccinia recondita. Canadian Journal of Plant Pathology* 5: 229–234.
- Eizenberg H, Plakhine D, Hershenhorn J, Kleifeld Y, Rubin B. 2003. Resistance to broomrape (*Orabanche* spp.) in sunflower (*Helianthus annuus* L.) is temperature dependent. *Journal of Experimental Botany* 54: 1305–1311.
- Ezuka A, Kaku H. 2000. A historical review of bacterial blight of rice. Bulletin of National Institute of Agrobiological Resources, Japan, 207.
- Fu D, Uauy C, Distelfeld A, Blechl A, Epstein L, Chen X, Sela H, Fahima T, Dubcovsky J. 2009. A kinase-START gene confers temperature-dependent resistance to wheat stripe rust. *Science* 323: 1357–1360.
- Garrett K, Dendy S, Frank E, Rouse M, Travers S. 2006. Climate change effects on plant disease: genomes to ecosystems. *Annual Review of Phytopathology* 44: 489–509.

Gibbs HL, Grant PR. 1987. Oscillating selection on Darwin's finches. *Nature* 327: 511–513.

Grant P, Grant B. 2002. Unpredictable evolution in a 30-year study of Darwin's Finches. *Science* 296: 707–711.

Guo A, Reimers P, Leach JE. 1993. Effect of light on incompatible interactions between *Xanthomonas oryzae* pv. *oryzae* and rice. *Physiological and Molecular Plant Pathology* 42: 413–425.

Hairston NG, Dillon TA. 1990. Fluctuating selection and response in a population of fresh-water copepods. *Evolution* 44: 1796–1805.

Horino O, Mew TW, Yamada T. 1982. The effect of temperature on the development of bacterial leaf blight on rice. *Annals of the Phytopathologi*cal Society of Japan 48: 72–75.

Iyer-Pascuzzi AS, Jiang H, Huang L, McCouch SR. 2008. Genetic and functional characterization of the rice bacterial blight disease resistance gene xa5. *Phytopathology* 98: 289–295.

Johnson R. 1984. A critical analysis of durable resistance. *Annual Review of Phytopathology* 22: 309–330.

Kauffman H, Reddy A, Hsiek S, Merca S. 1973. An improved technique for evaluating resistance of rice varieties to *Xanthomonas oryzae*. *Plant Disease Reporter* 57: 537–541.

Kolmer JA. 1996. Genetics of resistance to wheat leaf rust. Annual Review of Phytopathology 34: 435–455.

Leach JE, Vera Cruz CM, Bai JF, Leung H. 2001. Pathogen fitness penalty as a predictor of durability of disease resistance genes. *Annual Review of Phytopathology* 39: 187–224.

Leonard KJ, Czochor RJ. 1980. Theory of genetic interactions among populations of plants and their pathogens. *Annual Review of Phytopathol*ogy 18: 237–258.

Li ZK, Sanchez A, Angeles E, Singh S, Domingo J, Huang N, Khush G. 2001. Are the dominant and recessive plant disease resistance genes similar? A case study of rice R genes and *Xanthomonas oryzae* pv. oryzae races. *Genetics* 159: 757–765.

Lynch M. 1987. The consequences of fluctuating selection for isozyme polymorphisms in *Daphnia. Genetics* 115: 657–669.

Madden LV, Hughes G, van den Bosch F. 2007. The study of plant disease epidemics. St. Paul, MN, USA: APS Press.

Mazzola M, Leach JE, Nelson R, White FF. 1994. Analysis of the interaction between *Xanthomonas oryzae* pv. oryzae and the rice cultivars IR24 and IRB21. *Phytopathology* 84: 392–397.

Mew TW. 1987. Current status and future prospects of research on bacterial blight of rice. *Annual Review of Phytopathology* 25: 359–382.

Mew TW, Vera Cruz CM, Medalla ES. 1992. Changes in race frequency of *Xanthomonas oryzae* pv. *oryzae* in response to rice cultivars planted in the Philippines. *Plant Disease* 76: 1029–1032.

- Ogawa T, Yamamoto K, Khush G, Mew T. 1991. Breeding of near-isogenic lines of rice with single genes for resistance to bacterial blight pathogen (*Xanthomonas campestris* pv. *oryzae*). *Japan Journal of Breeding* 41: 523–529.
- Oña I, Vera Cruz CM, Nelson RJ, Leach JE, Mew TW. 1998. Epidemic development of bacterial blight on rice carrying resistance genes *Xa-4*, *Xa-7*, and *Xa-10. Plant Disease* 82: 1337–1340.

Ou SH. 1985. Rice diseases. Surrey, UK: Association Applied Biology.

Peng S, Huang J, Sheehy JE, Laza RC, Visperas RM, Zhong X, Centeno HGS, Khush G, Cassman KG. 2004. Rice yields decline with higher night temperature from global warming. *Proceedings of the National Academy of Sciences, USA* 101: 9971–9975.

Ponciano G, Webb KW, Bai JF, Vera Cruz CM, Leach JE. 2004. Molecular characterization of the *avrXa7* locus from *Xanthomonas oryzae* pv. *oryzae* field isolates. *Physiological and Molecular Plant Pathology* 64: 145–153.

Sheehy JE, Elmido A, Centeno HGS, Pablico P. 2005. Searching for new plants for climate change. *Journal of Agricultural Meteorology* 60: 463– 468.

Uauy C, Brevis JC, Chen X, Khan I, Jackson L, Chicaiza O, Distelfeld A, Fahima T, Dubcovsky J. 2005. High-temperature adult-plant (HTAP) stripe rust resistance gene Yr36 from *Triticum turgidum* ssp. dicoccoides is closely linked to the grain protein content locus Gpc-B1. *Theoretical* and Applied Genetics 112: 97–105.

Vera Cruz CM, Ardales EY, Skinner DZ, Talag J, Nelson RJ, Louws FJ, Leung H, Mew TW, Leach JE. 1996. Measurement of haplotypic variation in *Xanthomonas oryzae* pv. *oryzae* within a single field by rep-PCR and RFLP analyses. *Phytopathology* 86: 1352–1359.

Vera Cruz CM, Bai JF, Oña I, Leung H, Nelson RJ, Mew TW, Leach JE. 2000. Predicting durability of a disease resistance gene based on an assessment of the fitness loss and epidemiological consequences of avirulence gene mutation. *Proceedings of the National Academy of Sciences*, USA 97: 13500–13505.

Wang Z, Yamanouchi U, Katayose Y, Sasaki T, Yano M. 2001. Expression of the *Pib* rice-blast resistance gene family is up-regulated by environmental conditions favoring infection and by chemical signals that trigger secondary plant defenses. *Plant Molecular Biology* 47: 653–661.

Yamada T, Horino O, Samoto S. 1979. Studies on genetics and breeding of resistance to bacterial leaf blight in rice 1. Discovery of new varietal groups on the basis of reaction patterns to five different pathotypes of *Xanthomonas oryzae* (Uyeda et Ishiyama) Dowson in Japan. *Annals of the Phytopathological Society of Japan* 45: 240–246.